

# Cochrane Reviews für den Fachbereich Ergotherapie

Ressourcen zur Evidenzbasierung  
in den Gesundheitsfachberufen

Juli bis September 2018



Nutzerspezifische  
Cochrane Reviews



Die Cochrane Deutschland Stiftung analysiert monatlich alle [neu erschienenen Cochrane Reviews](#) nach Relevanz für die Gesundheitsfachberufe (GFB). Die Relevanz für die Disziplinen wird jeweils durch zwei Experten der GFB unabhängig voneinander beurteilt. Ebenso prüft die Cochrane Deutschland Stiftung, in wie weit die jeweiligen Cochrane Reviews für AWMF-Leitlinien relevant sind und ob sie dort zitiert werden.

Die Berichte können eine aktuelle und berufsspezifische Basis für Übersetzungsaktivitäten und andere Nutzungen von Cochrane Reviews in Forschung und Praxis werden. Für die Erarbeitung von Leitlinien können diese Übersichten ebenfalls hilfreich sein.

**Autoren:**

Katharina Wollmann & Sebastian Voigt-Radloff

**Kontakt:**

Cochrane Deutschland Stiftung  
Breisacher Str. 153  
D-79110 Freiburg  
[www.cochrane.de](http://www.cochrane.de)

Thieme H, Morkisch N, Mehrholz J, Pohl M, Behrens J, Borgetto B, Dohle C. Mirror therapy for improving motor function after stroke. Cochrane Database of Systematic Reviews 2018, Issue 7. Art. No.: CD008449. DOI: 10.1002/14651858.CD008449.pub3.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD008449.pub3/full/de#CD008449-abs-0004>

Publiziert 07/2018      Studien bis 2017

The results indicate evidence for the effectiveness of mirror therapy for improving upper extremity motor function, motor impairment, activities of daily living, and pain, at least as an adjunct to conventional rehabilitation for people after stroke. Major limitations are small sample sizes and lack of reporting of methodological details, resulting in uncertain evidence quality.

**Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Légaré F, Adekpedjou R, Stacey D, Turcotte S, Kryworuchko J, Graham ID, Lyddiatt A, Politi MC, Thomson R, Elwyn G, Donner-Banzhoff N. Interventions for increasing the use of shared decision making by healthcare professionals. Cochrane Database of Systematic Reviews 2018, Issue 7. Art. No.: CD006732. DOI: 10.1002/14651858.CD006732.pub4.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006732.pub4/full>

Publiziert 07/2018      Studien bis 2017

It is uncertain whether any interventions for increasing the use of SDM by healthcare professionals are effective because the certainty of the evidence is low or very low.

**Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Hopewell S, Adedire O, Copsey BJ, Boniface GJ, Sherrington C, Clemson L, Close JCT, Lamb SE. Multifactorial and multiple component interventions for preventing falls in older people living in the community. Cochrane Database of Systematic Reviews 2018, Issue 7. Art. No.: CD012221. DOI: 10.1002/14651858.CD012221.pub2.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012221.pub2/full#CD012221-sec1-0005>

Publiziert 07/2018      Studien bis 2017

Multifactorial interventions may reduce the rate of falls compared with usual care or attention control. However, there may be little or no effect on other fall-related outcomes. Multiple component interventions, usually including exercise, may reduce the rate of falls and risk of falling compared with usual care or attention control.

**Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Williams MA, Srikesavan C, Heine PJ, Bruce J, Brosseau L, Hoxey-Thomas N, Lamb SE. Exercise for rheumatoid arthritis of the hand. Cochrane Database of Systematic Reviews 2018, Issue 7. Art. No.: CD003832. DOI: 10.1002/14651858.CD003832.pub3.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD003832.pub3/full#CD003832-abs-0002>

Publiziert 07/2018      Studien bis 2017

It is uncertain whether exercise improves hand function or pain in the short term. It probably slightly improves function but has little or no difference on pain in the medium and long term. It is uncertain whether exercise improves grip and pinch strength in the short term, and probably has little or no difference in the medium and long term. The ACR50 response is unknown. People who received exercise with adherence strategies were probably more adherent in the medium term than who did not receive exercise, but with little or no difference in the long term. Hand exercise probably does not lead to adverse events. Future research should consider hand and wrist function as their primary outcome, describe exercise following the TIDieR guidelines, and evaluate behavioural strategies.

### **Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Cameron ID, Dyer SM, Panagoda CE, Murray GR, Hill KD, Cumming RG, Kerse N. Interventions for preventing falls in older people in care facilities and hospitals. Cochrane Database of Systematic Reviews 2018, Issue 9. Art. No.: CD005465. DOI: 10.1002/14651858.CD005465.pub4.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD005465.pub4/full#CD005465-sec1-0004>

Publiziert 09/2018      Studien bis 2017

In care facilities: we are uncertain of the effect of exercise on rate of falls and it may make little or no difference to the risk of falling. General medication review may make little or no difference to the rate of falls or risk of falling. Vitamin D supplementation probably reduces the rate of falls but not risk of falling. We are uncertain of the effect of multifactorial interventions on the rate of falls; they may make little or no difference to the risk of falling.

In hospitals: we are uncertain of the effect of additional physiotherapy on the rate of falls or whether it reduces the risk of falling. We are uncertain of the effect of providing bed sensor alarms on the rate of falls or risk of falling. Multifactorial interventions may reduce rate of falls, although subgroup analysis suggests this may apply mostly to a subacute setting; we are uncertain of the effect of these interventions on risk of falling.

### **Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Deshmukh SR, Holmes J, Cardno A. Art therapy for people with dementia. Cochrane Database of Systematic Reviews 2018, Issue 9. Art. No.: CD011073. DOI: 10.1002/14651858.CD011073.pub2.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD011073.pub2/full>

Publiziert 09/2018      Studien bis 2018

There is insufficient evidence about the efficacy of art therapy for people with dementia. More adequately-powered and high-quality studies using relevant outcome measures are needed.

### **Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Turner RR, Steed L, Quirk H, Greasley RU, Saxton JM, Taylor SJC, Rosario DJ, Thaha MA, Bourke L. Interventions for promoting habitual exercise in people living with and beyond cancer. Cochrane Database of Systematic Reviews 2018, Issue 9. Art. No.: CD010192. DOI: 10.1002/14651858.CD010192.pub3.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD010192.pub3/full>

Publiziert 09/2018      Studien bis 2018

Since the last version of this review, none of the new relevant studies have provided additional information to change the conclusions. We have found some improved understanding of how to encourage previously inactive cancer survivors to achieve international physical activity guidelines. Goal setting, setting of graded tasks and instruction of how to perform behaviour, feature in interventions that meet recommendations targets and report adherence of 75% or more. However, long-term follow-up data are still limited, and the majority of studies are in white women with breast cancer. There are still a considerable number of published studies with numerous and varied issues related to high risk of bias and poor reporting standards. Additionally, the meta-analyses were often graded as consisting of low- to very low-certainty evidence. A very small number of serious adverse effects were reported amongst the studies, providing reassurance exercise is safe for this population.

### **Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT

Yamamoto S, Hotta K, Ota E, Matsunaga A, Mori R. Exercise-based cardiac rehabilitation for people with implantable ventricular assist devices. Cochrane Database of Systematic Reviews 2018, Issue 9. Art. No.: CD012222. DOI: 10.1002/14651858.CD012222.pub2.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012222.pub2/full>

Publiziert 09/2018      Studien bis 2017

The evidence is currently inadequate to assess the safety and efficacy of exercise-based CR for people with implantable VADs compared with usual care. The amount of RCT evidence was very limited and of very low quality. In addition, the training duration was very short term, that is from six to eight weeks. Further high-quality and well-reported RCTs of exercise-based CR for people with implantable VADs are needed. Such trials need to collect data on events (mortality and rehospitalisation), patient-related outcomes (including quality of life), and cost-effectiveness.

### **Relevante AWMF-Leitlinien, die das Cochrane Review enthalten (CR IN) bzw. nicht enthalten (CR OUT)**

CR IN

CR OUT